

CSC165 fall 2014

Mathematical expression

Danny Heap

heap@cs.toronto.edu

BA4270 (behind elevators)

<http://www.cdf.toronto.edu/~heap/165/F14/>

416-978-5899

Course notes, chapter 4



Outline

Bounded below

some theorems

notes

annotated slides



how to prove $n^3 \notin \mathcal{O}(3n^2)$?

Negate $\exists c \in \mathbb{R}^+, \exists B \in \mathbb{N}, \forall n \in \mathbb{N}, n \geq B \Rightarrow n^3 \leq c3n^2\}$



non-polynomials

Big-oh statements about polynomials are pretty easy to prove:
 $f \in \mathcal{O}(g)$ exactly when the highest-degree term of g is no smaller than the highest-degree term of f .

What about functions such as $\log(n)$ or 3^n ? Logarithmic functions are in big-Oh of **any** polynomial, whereas exponential functions (with a base bigger than 1) are not in big-Oh of any polynomial. How do you prove such things?

Prove $2^n \notin \mathcal{O}(n^2)$

Use $\lim_{n \rightarrow \infty} 2^n / n^2$

Do you know anything about the ratio $2^n / n^2$, as n gets very large? How do you evaluate:

$$\lim_{n \rightarrow \infty} \frac{2^n}{n^2}$$

If the limit evaluates to ∞ , then that's the same as saying:

$$\forall c \in \mathbb{R}^+, \exists n_c \in \mathbb{N}, \forall n \in \mathbb{N}, n \geq n_c \Rightarrow \frac{2^n}{n^2} > c$$

Once your enemy hands you a c , you can choose an n_c with the required property.



prove $2^n \notin \mathcal{O}(n^2)$

Using l'Hôpital's rule and limits



big-Omega

Notice that the definition of big-Omega differs in just one character from big-Oh:

$$\Omega(g) = \{f : \mathbb{N} \mapsto \mathbb{R}^{\geq 0} \mid \exists c \in \mathbb{R}^+, \exists B \in \mathbb{N}, \forall n \in \mathbb{N}, n \geq B \Rightarrow f(n) \geq cg(n)\}$$

The rôle of B is, as with big-Oh, to act as a breakpoint, so comparisons don't have to start at the origin.

The rôle of c is to scale g *down* below f .

If you're proving $f \in \Omega(g)$, you get to choose c and B to suit your proof. Notice that it would be really unfair to allow c to be zero.

one more bound

It often happens that functions are bounded above *and* below by the same function. In other words, $f \in \mathcal{O}(G) \wedge f \in \Omega(g)$. We combine these two concepts into $f \in \Theta(g)$.

$$\Theta(g) = \{f : \mathbb{N} \mapsto \mathbb{R}^{\geq 0} \mid \exists c_1 \in \mathbb{R}^+, \exists c_2 \in \mathbb{R}^+, \exists B \in \mathbb{N}, \\ \forall n \in \mathbb{N}, n \geq B \Rightarrow c_1 g(n) \leq f(n) \leq c_2 g(n)\}$$

You might want to draw pictures, and conjecture about appropriate values of c_1, c_2, B for $f = 5n^2 + 15$ and $g = n^2$.

How to prove general statement about two functions:

$$\mathcal{F} = \{f : \mathbb{N} \mapsto \mathbb{R}^{\geq 0}\}$$

$$\forall f, g, h \in \mathcal{F}, (f \in \mathcal{O}(g) \wedge g \in \mathcal{O}(h)) \Rightarrow f \in \mathcal{O}(h)$$



how about:

$$\mathcal{F} = \{f : \mathbb{N} \mapsto \mathbb{R}^{\geq 0}\}$$

$$\forall f, g \in \mathcal{F}, f \in \mathcal{O}(g) \Rightarrow g \in \Omega(f)$$



prove or disprove:

$$\mathcal{F} = \{f : \mathbb{N} \mapsto \mathbb{R}^{\geq 0}\}$$

$$\forall f, g \in \mathcal{F}, f \in \mathcal{O}(g) \Rightarrow f \cdot f \in \mathcal{O}(g \cdot g)$$



prove or disprove:

$$\mathcal{F} = \{f : \mathbb{N} \mapsto \mathbb{R}^{\geq 0}\}$$

$$\forall f, g \in \mathcal{F}, (f \in \mathcal{O}(h) \wedge g \in \mathcal{O}(h)) \Rightarrow (f + g) \in \mathcal{O}(h)$$



prove or disprove:

$$\mathcal{F} = \{f : \mathbb{N} \mapsto \mathbb{R}^{\geq 0}\}$$

$$\forall f, g \in \mathcal{F}, f \in \mathcal{O}(g) \Rightarrow f \in \mathcal{O}(g \cdot g)$$



prove or disprove:

$$\mathcal{F} = \{f : \mathbb{N} \mapsto \mathbb{R}^{\geq 0}\}$$

$$\forall f, g, h \in \mathcal{F}, (f \in \Omega(g) \wedge f \in \Omega(h)) \Rightarrow f \in \Omega(g + h)$$



Notes

annotated slides

- ▶ monday's annotated slides
- ▶ wednesday's annotated slides
- ▶ friday's annotated slides

