

PLEASE HAND IN

UNIVERSITY OF TORONTO
Faculty of Arts and Science

Term test #2

CSC 165H1

Duration — 50 minutes

PLEASE HAND IN

aids allowed: 8.5" x 11" handwritten aid sheet, both sides

Last Name: _____

First Name: _____

Do not turn this page until you have received the signal to start.
(In the meantime, please fill out the identification section above,
and read the instructions below.)

This test consists of 3 questions on 4 pages (including this one). *When you receive the signal to start, please make sure that your copy of the test is complete.*

Please answer questions in the space provided. You will earn 20% for any question you leave blank or write "I cannot answer this question," on.

Good Luck!

Question 1. [10 MARKS]

Use the proof structure from this course (including comments) to **prove**:

$$\forall m \in \mathbb{N}, \forall n \in \mathbb{N}, (\exists i \in \mathbb{N}, m = 5i + 1) \wedge (\exists j \in \mathbb{N}, n = 5j + 2) \Rightarrow (\exists k \in \mathbb{N}, m + n = 5k + 3)$$

Proof:

Assume $m \in \mathbb{N}, n \in \mathbb{N}$ # generic natural numbers

Assume $(\exists i \in \mathbb{N}, m = 5i + 1) \wedge (\exists j \in \mathbb{N}, n = 5j + 2)$ # the antecedent

Let $i_0 \in \mathbb{N}$ be such that $m = 5i_0 + 1$

Let $j_0 \in \mathbb{N}$ be such that $n = 5j_0 + 2$

then $m + n = (5i_0 + 1) + (5j_0 + 2) = 5(i_0 + j_0) + 3$

Let $k = i_0 + j_0$

then $k \in \mathbb{N}$ # i_0 and j_0 are both natural numbers

then $m + n = 5k + 3$

then $\exists k \in \mathbb{N}, m + n = 5k + 3$

then $(\exists i \in \mathbb{N}, m = 5i + 1) \wedge (\exists j \in \mathbb{N}, n = 5j + 2) \Rightarrow \exists k \in \mathbb{N}, m + n = 5k + 3$

then $\forall m \in \mathbb{N}, \forall n \in \mathbb{N}, (\exists i \in \mathbb{N}, m = 5i + 1) \wedge (\exists j \in \mathbb{N}, n = 5j + 2) \Rightarrow (\exists k \in \mathbb{N}, m + n = 5k + 3)$

Question 2. [10 MARKS]

For $x \in \mathbb{R}$, define $\lfloor x \rfloor$ by

$$\lfloor x \rfloor \in \mathbb{Z} \wedge \lfloor x \rfloor \leq x \wedge (\forall z \in \mathbb{Z}, z \leq x \Rightarrow z \leq \lfloor x \rfloor)$$

Use this definition of $\lfloor x \rfloor$, and the structured proof technique from this course (including comments, where appropriate), to **prove**:

$$\forall e \in \mathbb{R}^+, \exists d \in \mathbb{R}^+, \forall w \in \mathbb{R}, w > d \Rightarrow \lfloor w \rfloor > e$$

Proof:

Assume $e \in \mathbb{R}^+$ # generic positive real number

Pick $d = e + 1$, then $d \in \mathbb{R}^+$

Assume $w \in \mathbb{R}$ # generic real number

Assume $w > d$ # the antecedent

then $w > e + 1$ # because of the d we picked

Assume $\lfloor w \rfloor \leq e$ # for contradiction

then $\lfloor w \rfloor + 1 \leq e + 1$ # add 1 to both side

then $\lfloor w \rfloor + 1 < w$ # transitivity

and $\lfloor w \rfloor + 1 \in \mathbb{Z}$ # $\lfloor w \rfloor$ and 1 are both integers

then $\lfloor w \rfloor + 1 \leq \lfloor w \rfloor$ # by definition of $\lfloor w \rfloor$

then $1 \leq 0$ # subtract $\lfloor w \rfloor$ from both sides, and contradict with $1 > 0$

then $\lfloor w \rfloor > e$ # because of contradiction

then $w > d \Rightarrow \lfloor w \rfloor > e$ # introduce \Rightarrow

then $\forall w \in \mathbb{R}, w > d \Rightarrow \lfloor w \rfloor > e$ # introduce \forall

then $\exists d \in \mathbb{R}^+, \forall w \in \mathbb{R}, w > d \Rightarrow \lfloor w \rfloor > e$ # introduce \exists

then $\forall e \in \mathbb{R}^+, \exists d \in \mathbb{R}^+, \forall w \in \mathbb{R}, w > d \Rightarrow \lfloor w \rfloor > e$ # introduce \forall

Alternative Proof utilising the result $\lfloor w \rfloor > w - 1$

Assume $e \in \mathbb{R}^+$ # generic positive real number

Pick $d = e + 1$, then $d \in \mathbb{R}^+$

Assume $w \in \mathbb{R}$ # generic real number

Assume $w > d$ # the antecedent

then $w > e + 1$ # because of the d we picked

then $w - 1 > e$ # subtract 1 from both sides

and $\lfloor w \rfloor > w - 1$ # property proven in lecture

then $\lfloor w \rfloor > e$ # transitivity

then $w > d \Rightarrow \lfloor w \rfloor > e$ # introduce \Rightarrow

then $\forall w \in \mathbb{R}, w > d \Rightarrow \lfloor w \rfloor > e$ # introduce \forall

then $\exists d \in \mathbb{R}^+, \forall w \in \mathbb{R}, w > d \Rightarrow \lfloor w \rfloor > e$ # introduce \exists

then $\forall e \in \mathbb{R}^+, \exists d \in \mathbb{R}^+, \forall w \in \mathbb{R}, w > d \Rightarrow \lfloor w \rfloor > e$ # introduce \forall

Question 3. [10 MARKS]

For $x \in \mathbb{R}$, define $\lfloor x \rfloor$ by

$$\lfloor x \rfloor \in \mathbb{Z} \wedge \lfloor x \rfloor \leq x \wedge (\forall z \in \mathbb{Z}, z \leq x \Rightarrow z \leq \lfloor x \rfloor)$$

Use this definition of $\lfloor x \rfloor$, and the proof structure from this course (including comments) to **disprove**:

$$\exists z \in \mathbb{Z}, \exists x \in \mathbb{R}, \lfloor x \rfloor + z \neq \lfloor x + z \rfloor$$

Sample solution: First negate the statement, yielding

$$\forall z \in \mathbb{Z}, \forall x \in \mathbb{R}, \lfloor x \rfloor + z = \lfloor x + z \rfloor$$

Like what we did in the assignment, we prove this equality by showing

$$(\lfloor x \rfloor + z \leq \lfloor x + z \rfloor) \wedge (\lfloor x + z \rfloor \leq \lfloor x \rfloor + z)$$

Proof:

Assume $z \in \mathbb{Z}$ # generic integer

Assume $x \in \mathbb{R}$ # generic real number

then $\lfloor x \rfloor \leq x$ # by definition of $\lfloor x \rfloor$

then $\lfloor x \rfloor + z \leq x + z$ # add x to both sides

and $\lfloor x \rfloor + z \in \mathbb{Z}$ # both $\lfloor x \rfloor$ and z are integers

then $\lfloor x \rfloor + z \leq \lfloor x + z \rfloor$ # definition of $\lfloor x + z \rfloor$, first inequality obtained

and $\lfloor x + z \rfloor \leq x + z$ # definition of $\lfloor x + z \rfloor$

then $\lfloor x + z \rfloor - z \leq x$ # subtract z from both sides

and $\lfloor x + z \rfloor - z \in \mathbb{Z}$ # both $\lfloor x + z \rfloor$ and z are integers

then $\lfloor x + z \rfloor - z \leq \lfloor x \rfloor$ # definition of $\lfloor x \rfloor$

then $\lfloor x + z \rfloor \leq \lfloor x \rfloor + z$ # add z to both sides, second inequality obtained

then $(\lfloor x \rfloor + z \leq \lfloor x + z \rfloor) \wedge (\lfloor x + z \rfloor \leq \lfloor x \rfloor + z)$ # conjunction introduction

then $\lfloor x \rfloor + z = \lfloor x + z \rfloor$ # $a = b \Leftrightarrow a \leq b \wedge b \leq a$

then $\forall x \in \mathbb{R}, \lfloor x \rfloor + z = \lfloor x + z \rfloor$ # introduce \forall

then $\forall z \in \mathbb{Z}, \forall x \in \mathbb{R}, \lfloor x \rfloor + z = \lfloor x + z \rfloor$ # introduce \forall

1: _____/10

2: _____/10

3: _____/10

TOTAL: _____/30